

# Potential Designated Chemicals

# Selected Aroma Chemicals

Gail Krowech, Laurel Plummer, and Sara Hoover  
Office of Environmental Health Hazard Assessment

*Presentation to Scientific Guidance Panel  
Sacramento, CA*

*November 14, 2013*



# What are designated chemicals?

- ▶ Chemicals that can be considered for biomonitoring by the Program
- ▶ Consist of
  - Chemicals that are part of CDC's National Reports on Human Exposure to Environmental Chemicals program
  - Chemicals that the Scientific Guidance Panel has recommended be added to the list of designated chemicals

# Background


November 2012 SGP Meeting:

- ▶ Presentation on screening of four classes of synthetic musks and a structurally related aroma chemical (Iso E Super<sup>®</sup>)
- ▶ SGP requested documents to support consideration of these aroma chemicals as potential designated chemicals


# Implementing SGP recommendation

- ▶ Two classes for consideration today – structurally similar, common analytical method
  - Synthetic polycyclic musks
  - Tetramethyl acetyloctahydronaphthalenes
- ▶ Other classes not under consideration today –
  - Nitro musks – low or no current use
  - For future consideration
    - Macrocyclic musks
    - Alicyclic musks

# Polycyclic musks – example structures



**HHCB\***  
Galaxolide®




**AHTN\*\***  
Tonalide®

\*HHCB: 1,3,4,6,7,8-Hexahydro-4,6,6,7,8,8-hexamethylcyclopenta[*g*]-2-benzopyran


\*\*AHTN: 7-Acetyl-1,1,3,4,4,6-hexamethyltetrahydronaphthalene

# Tetramethyl acetyloctahydronaphthalenes



**OTNE\***  
(beta isomer, Iso-E Super®)

OTNE is structurally similar to  
some polycyclic musks



**AHTN**


\*OTNE: 1-(1,2,3,4,5,6,7,8-Octahydro-2,3,8,8-tetramethyl-2-naphthalenyl)ethanone

# Criteria for Panel to recommend designated chemicals


- ▶ **Exposure or potential exposure to the public or specific subgroups**
- ▶ **The known or suspected health effects based on peer-reviewed scientific studies**
- ▶ **The need to assess the efficacy of public health actions to reduce exposure**
- ▶ **The availability of a biomonitoring analytical method with adequate accuracy, precision, sensitivity, specificity, and speed**
- ▶ **The availability of adequate biospecimen samples**
- ▶ **The incremental analytical cost to perform the biomonitoring analysis for the chemical**

# Polycyclic musks

- ▶ Widely used in personal care products and some cleaning products
- ▶ Replacements for nitro musks (e.g., musk xylene)
- ▶ Highlighting two – HHCB and AHTN – which have been commercially most important




HHCB




AHTN


# Other polycyclic musks




**ADBI<sup>1</sup>**




**ATII<sup>3</sup>**



**AHMI<sup>2</sup>**



**DPMI<sup>4</sup>**



**AETT<sup>5</sup>**

<sup>1</sup> ADBI: (4-Acetyl-1,1-dimethyl-6-*tert*-butylindan)

<sup>2</sup> AHMI: 6-Acetyl-1,1,2,3,3,5-hexamethylindane

<sup>3</sup> ATII : 5-Acetyl-1,1,2,6-tetramethyl-3-isopropylindan

<sup>4</sup> DPMI: 6,7-Dihydro-1,1,2,3,3-pentamethyl-4[5*H*]indanon

<sup>5</sup> AETT: Acetyethyltetramethyltetralin

# Polycyclic musks

## U.S. Production/Import Volume (pounds)

	1986	1994	1998	2002	2006	2012
HHCB	500K-1M	1-10M	1-10M	1-10M	1-10M	3.1 M
AHTN	10-500K	10-500K	1-10M	NR	NR	CBI <i>*220-330K</i>
DPMI	10-500K	10-500K	10-500K	10-500K	NR	CBI

Table notes:

Source: U.S. EPA (2002;2006; 2012) unless otherwise noted

NR=not reported; volume is less than U.S. EPA reporting threshold

CBI= Reported as Confidential Business Information

\*AHTN volume of use in North America, reported as 220-330K lbs in 2011 (IFRA-NA)

# *Polycyclic musks*

## Use and exposure

### Personal care products

- Perfumes / fragrances
- Body lotions / body creams
- Deodorants / antiperspirants
- Shower gels / shaving cream
- Shampoo / conditioner products
- Hand soaps / bar soaps

Reiner and Kannan (2006)  
Dodson et al. (2012)

# *Polycyclic musks*

## Use and exposure

### Household products

- Carpet cleaner
- Furniture polish
- Dish soap
- Laundry detergent
- Stain remover
- Fabric softener
- Liquid bleach
- Disinfecting wipes

Reiner and Kannan (2006)  
Dodson et al. (2012)

## *Polycyclic musks:*

# Example levels in personal care products

### ▶ HHCB

- Body splash 4,990  $\mu\text{g/g}$
- Body lotion 3,740  $\mu\text{g/g}$
- Deodorant 2,250  $\mu\text{g/g}$
- Shaving cream 1,230  $\mu\text{g/g}$

### ▶ AHTN

- Perfume 451  $\mu\text{g/g}$
- Deodorant 438  $\mu\text{g/g}$
- Body cream 145  $\mu\text{g/g}$

# *Polycyclic musks:*

## Example levels in consumer products

### *Personal care products*

- Bar soap >100–1000 µg/g (HHCB); >1–100 µg/g (AHTN)
- Hand soap >1–100 µg/g (HHCB, DPMI)

### *Household cleaning products*

- Dish liquid >100–1000 µg/g (HHCB)
- Carpet cleaner >100–1000 µg/g (HHCB)
- Laundry detergent >1–100 µg/g (HHCB, AHTN)
- Dryer sheets >1–100 µg/g (HHCB, AHTN)
- Polish/wax >1–100 µg/g (HHCB, DPMI)
- Air freshener >1–100 µg/g (HHCB)

# *Polycyclic musks*

## Levels in house dust

Samples collected as part of the Canadian House Dust Study, 2007–2010 (n=49)

Household vacuum cleaner dust			
	Detection frequency (%)	Median (ng/g)	Range (ng/g)
HHCB	100	992	36–31,100
AHTN	100	405	91–2,360

Kubwabo et al. (2012)

# *Polycyclic musks*

## Environmental occurrence in U.S.

- ▶ Main environmental source is effluent from wastewater treatment plants (WWTPs)
  - HHCB and AHTN detected in:
    - fish caught in WWTP effluent waters (sampled in 2006)
    - sewage sludge (biosolids)
    - some drinking water
    - run-off from agricultural fields irrigated with treated wastewater (California)

# *Polycyclic musks*

## Detections in biota

### ▶ Bivalves in San Francisco Bay

- HHCB, AHTN, ADBI, AETT detected in 2002–2003 sampling
- HHCB, AHTN, ADBI, AETT detected in mussels in 2009–2010 sampling

### ▶ Fish

- Levels dependent on location, and on metabolism and lipid content of fish

### ▶ Marine mammals

- Finless porpoises (Japan): Level in one porpoise was comparable to level in its fetus

# *Polycyclic musks*

## Known or suspected health effects

- ▶ Indications of endocrine activity
  - *In vitro*
    - Weak estrogenicity
    - Inhibition of estrogen, androgen, and progesterone activity
    - Decreased progesterone and cortisol synthesis
  - *In vivo*
    - Anti-estrogenicity (transgenic zebrafish, trout)
  
- ▶ Other *in vitro* biological activity
  - AHTN caused changes in the activation of certain signaling pathways (mouse embryonic stem cells)
  - Several polycyclic musks inhibited efflux transporters (mussel gill tissue)

# Properties of polycyclic musks

- ▶ Lipophilic chemicals

Polycyclic musk	Log $K_{ow}$
HHCB	5.9*
AHTN	5.7*
ADBI	5.4**
AHMI	5.8**
DPMI	4.5**
AETT	6.4 ( <i>est</i> )*

- ▶ Potential to bioaccumulate in some species
- ▶ Some indications of persistence (e.g., experimental studies in soils amended with sludge)

\*SRC (2013)

\*\*Cited in Rimkus et al. (1999)

# *Polycyclic musks*

## Biomonitoring studies

- ▶ Multiple studies in blood, breast milk, adipose tissue (HHCB and AHTN)
- ▶ Several studies reported levels increased with use of personal care products
- ▶ Most studies from Europe and Asia
- ▶ Few studies from the U.S.

# *Polycyclic musks*


## Biomonitoring studies

Breast milk – Massachusetts (n=39)

- ▶ HHCb: detection frequency: 97%
  - Mean: 220 ng/g lipid
  - Range: <5 – 917 ng/g lipid
- ▶ AHTN: detection frequency: 56%
  - Mean: 46.8 ng/g lipid
  - Range: <5–144 ng/g lipid

Reiner et al. (2007)

# Tetramethyl acetyloctahydronaphthalenes



OTNE\*

(beta isomer, Iso-E Super®)

- ▶ Woody, floral, or amber fragrances
- ▶ Widely used in personal care products and some cleaning products

\*OTNE: 1-(1,2,3,4,5,6,7,8-Octahydro-2,3,8,8-tetramethyl-2-naphthalenyl)ethanone

# Tetramethyl acetyloctahydronaphthalenes

## U.S. Production/Import Volume (pounds)

	1986	1994	1998	2002	2006	2012
54464-57-2 (beta isomer)	10K-500K	500K-1M	500K-1M	1M-10M	1M-10M	1M-10M
68155-67-9 (alpha isomer)	NR	10K-500K	500K-1M	1M-10M	1M-10M	CBI
68155-66-8 (gamma isomer)	NR	10K-500K	500K-1M	500K-1M	1M-10M	CBI
54464-59-4 ("4 <sup>th</sup> " isomer)	NR	NR	NR	NR	500K-1M	CBI

Source: U.S. EPA (2002;2006;2012)

NR=not reported; volume less than U.S. EPA reporting threshold

CBI= Reported as Confidential Business Information

# Uses and exposure: Examples

- ▶ Personal care products
  - Perfume/cologne
  - Soap/shower gels/shampoo
  - Body lotion/skin conditioner
  
- ▶ Cleaning products
  - Air freshener
  - Laundry detergent
  - Fabric softener

# Levels of OTNE in house dust

Samples collected as part of the Canadian House Dust Study, 2007–2010 (n=49)

Household vacuum cleaner dust			
	Detection frequency (%)	Median (ng/g)	Range (ng/g)
OTNE	82	212	nd – 5,620
<i>Compared to polycyclic musks</i>			
HHCB	100	992	36 – 31,100
AHTN	100	405	91 – 2,360

nd = not detected

Kubwabo et al. (2012)

# *Tetramethyl acetyloctahydronaphthalenes*

## Environmental occurrence

- ▶ Main environmental source is effluent from wastewater treatment plants (WWTPs)
- ▶ OTNE detected in:
  - Influent and effluent wastewater
  - Sewage sludge
    - Levels comparable to the polycyclic musks HHCB and AHTN

# *Tetramethyl acetyloctahydronaphthalenes*

## Bioaccumulation and persistence

- ▶ Bioaccumulation
  - Lipophilic:  $\log K_{ow} > 5$
  - Experimental BCFs (Bioconcentration Factors) do not suggest bioaccumulation (below 1000)
- ▶ Persistence
  - Few published studies
  - No evidence of persistence based on available data

# *Tetramethyl acetyloctahydronaphthalenes*

## Known or suspected health effects

- ▶ Few toxicological data for tetramethyl acetyloctahydronaphthalenes are publicly available
- ▶ Structurally similar to AHTN, which has shown some potential for endocrine and other biological activity

# Summary – Polycyclic musks

- ▶ High levels in personal care and household cleaning products
- ▶ Potential to bioaccumulate in some species
- ▶ Potential for endocrine and other biological activity
- ▶ Detected in:
  - Various environmental samples, including house dust
  - Human blood, breast milk, adipose tissue samples

# Summary – Tetramethyl acetyloctahydronaphthalenes

- ▶ OTNE – high production volume chemical
- ▶ Detected in dust, wastewater treatment plant influent and effluent, biosolids
- ▶ Structurally similar to AHTN

# Laboratory analysis

- ▶ Methods for analysis of some of these chemicals available in the literature
- ▶ Laboratory would develop methods to measure polycyclic musks and tetramethyl acetyloctahydronaphthalenes in serum samples
- ▶ Analysis could likely be bundled

# Need to assess efficacy of public health actions

- ▶ Widespread use of these aroma chemicals in California and in the U.S.
- ▶ Biomonitoring would:
  - Determine whether these chemicals are found in California residents and at what levels
  - Track levels over time

# Options for the Panel

- ▶ Designate: “synthetic polycyclic musks” as a class
- ▶ Designate: “tetramethyl acetyloctahydronaphthalenes” as a class
- ▶ Postpone decision
- ▶ Decide against designating